The Polarizations of Gravitational Waves in Scalar-Tensor Theory

Shaoqi Hou,* Yungui Gong,† and Yunqi Liu‡

School of Physics, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China

arXiv:1704.01899v2 [gr-qc] 13 Apr 2017

Abstract

We analyze the polarization states of gravitational waves in Horndeski theory. In addition to the familiar plus and cross modes appearing in Einstein's general relativity, there is one more polarization state which is the mixture of the transverse breathing and longitudinal modes. The additional mode is excited by the massive scalar field. In the massless limit, the longitudinal mode disappears, while the breathing one persists. The upper bound on the graviton mass severely constrains the amplitude of the longitudinal mode, which makes its detection highly unlikely by the ground-based or space-borne interferometers in the near future. However, pulsar timing arrays might be able to observe the longitudinal mode.

PACS numbers: 04.30.Nk, 04.50.Kd

* shou1397@hust.edu.cn

† yggong@hust.edu.cn

‡ liuyunqi@hust.edu.cn

I. INTRODUCTION

The first detection of gravitational waves by the Laser Interferometer Gravitational-Wave Observatory (LIGO) further supports Einstein's general relativity (GR) and provides a new tool to study gravitational physics [1]. To confirm gravitational waves in GR, we need to determine the polarizations of gravitational waves and this will be done by the network of ground-based Advanced LIGO (aLIGO) and Virgo or the future space-borne Laser Interferometer Space Antenna (LISA). In GR, the gravitational wave propagates at the speed of light and it has two polarization states, the plus and cross modes. In alternative metric theories of gravity, there may exist six polarization states. For null plane gravitational waves which propagate at the speed of light, the six polarization states are classified by the little group $E(2)$ of the Lorentz group with the help of the six independent Newman-Penrose variables Ψ_2, Ψ_3, Ψ_4 and Φ_{22} [2–4]. In particular, the complex variable Ψ_4 denotes the familiar plus and cross modes in GR. In Brans-Dicke theory [5], in addition to the plus and cross modes Ψ_4 of the massless gravitons, there exists another breathing mode Φ_{22} due to the massless Brans-Dicke scalar field [3].

Brans-Dicke theory is a simple extension to GR. In Brans-Dicke theory, gravitational interaction is mediated by both the metric tensor and the Brans-Dicke scalar field, and the Brans-Dicke field takes the role of Newton's gravitational constant. In more general scalar-tensor theories of gravity, the scalar field has self-interaction and it is usually massive. In 1974, Horndeski found the most general scalar-tensor theory of gravity whose equation of motion is at most the second order differential equation [6]. Even though there are higher derivative terms, there is no Ostrogradsky instability [7] and no more degree of freedom is introduced in Horndeski theory, so we expect that there is an extra polarization state in addition to the plus and cross modes. If the scalar field is massless, then the additional polarization state should be the breathing mode Φ_{22}.

When the interaction between quantized matter fields and the classical gravitational field is considered, the quadratic terms $R_{\mu\nu\alpha\beta}R^{\mu\nu\alpha\beta}$ and R^2 are needed as counter terms to remove the singularities in the energy-momentum tensor [8]. Even though the quadratic gravitational theory is renormalizable [9], the theory has ghost because of the presence of higher derivatives [9, 10]. However, the general nonlinear $f(R)$ gravity [11] has no ghost and it is equivalent to a scalar-tensor theory of gravity [12, 13]. The effective mass of

the equivalent scalar field is $f'(0)/3f''(0)$, and the massive scalar field generates both the longitudinal and transverse breathing modes [14–16]. The polarizations of gravitational waves in $f(R)$ gravity were also discussed in [17–23]. The authors in [19, 20] calculated the Newman-Penrose variables and found that Ψ_2, Ψ_4 and Φ_{22} are nonzero. They then claimed that there are four polarization states in $f(R)$ gravity. Recently, it was pointed out that the direct application of the Newman-Penrose formalism derived for the null plane gravitational waves to the massive field is not correct, and the polarization state of the massive field is the mixture of the longitudinal and the transverse breathing modes [23]. Furthermore, the longitudinal force is independent of the effective mass of the equivalent scalar field, so it cannot be used to understand how the polarization reduces to the transverse breathing mode in the massless limit.

In this paper, the focus is on the polarizations of gravitational waves in the most general scalar-tensor theory, the Horndeski theory, and the possible constraints on the theory [24, 25]. It is assumed that matter minimally couples to the metric, so test particles follow geodesics. The gravitational wave solutions are obtained from the linearized equations of motion around the flat space-time background, and the geodesic deviation equations are used to reveal the polarization modes of the massive scalar field. The analysis shows that in Horndeski theory, the massive scalar field excites both breathing and longitudinal polarizations. The effect of the longitudinal polarization on the geodesic deviation depends on the mass and it is smaller than that of the transverse polarization in the aLIGO and LISA frequency bands. In the massless limit, the longitudinal mode disappears, while the breathing mode persists. As a special example of the scalar-tensor theory, the result is applied to discuss $f(R)$ gravity. The paper is organized as follows. In Section II, the wave equation and the polarization states of gravitational waves in Horndeski theory are considered. We conclude the paper in Section III.

II. GRAVITATIONAL WAVES IN HORNDESKI GRAVITY

The most general scalar-tensor theory whose equation of motion is at most the second order differential equation is the Horndeski theory with the the following action [6],

$$S = \int d^4x\sqrt{-g}(L_2 + L_3 + L_4 + L_5), \tag{1}$$

where

$$
\begin{aligned}
L_2 = K(\phi, X), \quad L_3 = -G_3(\phi, X)\Box\phi, \\
L_4 = G_4(\phi, X)R + G_{4,X}\left[(\Box\phi)^2 - (\nabla_\mu\nabla_\nu\phi)(\nabla^\mu\nabla^\nu\phi)\right], \\
L_5 = & G_5(\phi, X)G_{\mu\nu}\nabla^\mu\nabla^\nu\phi - \frac{1}{6}G_{5,X}\left[(\Box\phi)^3\right. \\
& - 3(\Box\phi)(\nabla_\mu\nabla_\nu\phi)(\nabla^\mu\nabla^\nu\phi) \\
& \left. + 2(\nabla^\mu\nabla_\alpha\phi)(\nabla^\alpha\nabla_\beta\phi)(\nabla^\beta\nabla_\mu\phi)\right],
\end{aligned} \tag{2}
$$

$X = -\nabla_\mu\phi\nabla^\mu\phi/2$, $\Box\phi = \nabla_\alpha\nabla^\alpha\phi$, the functions K, G_3, G_4 and G_5 are arbitrary functions of ϕ and X, and $G_{4,X}(\phi, X) = dG_4(\phi, X)/dX$. For Einstein's general relativity, $K = G_3 = G_5 = 0$ and $G_4 = 1/(16\pi G)$. For Brans-Dicke theory, $G_3 = G_5 = 0$, $K = \omega X/\phi$ and $G_4 = \phi$. For general scalar-tensor theory of gravity, $G_3 = G_5 = 0$, $K = \omega(\phi)X - V(\phi)$ and $G_4 = F(\phi)$. Since $f(R)$ gravity can be written as the scalar-tensor theory of gravity, so for $f(R)$ gravity, we have $G_3 = G_5 = 0$, $K = f(\phi) - \phi f'(\phi)$ and $G_4 = f'(\phi)$ with $f'(\phi) = df(\phi)/d\phi$. To get the flat background solution $g_{\mu\nu} = \eta_{\mu\nu}$ and $\phi = \phi_0$ with a constant ϕ_0, we require $K(\phi_0, 0) = 0$ and $K_{,\phi_0} = dK(\phi, X)/d\phi|_{\phi=\phi_0, X=0}$. Now we expand the fields around the background, $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ and $\phi = \phi_0 + \varphi$. To the first order of approximation, we get the linearized equations of motion,

$$
(\Box - m^2)\varphi = 0, \tag{3}
$$

$$
G^{(1)}_{\mu\nu} - \frac{G_{4,\phi_0}}{G_4(0)}(\partial_\mu\partial_\nu\varphi - \eta_{\mu\nu}\Box\varphi) = 0, \tag{4}
$$

where $G^{(1)}_{\mu\nu}$ is the linearized Einstein tensor, $G_4(0) = G_4(\phi_0, 0)$, $K_{,X_0} = dK(\phi, X)/dX|_{\phi=\phi_0, X=0}$ and the mass of the scalar field is

$$
m^2 = -\frac{K_{,\phi_0\phi_0}}{K_{,X_0} - 2G_{3,\phi_0} + 3G^2_{4,\phi_0}/G_4(0)}. \tag{5}
$$

For Brans-Dicke theory, $K_{,\phi_0\phi_0} = 0$, so $m^2 = 0$. For the general scalar-tensor theory with $G_4 = F(\phi)$ and $K = \omega(\phi)X - V(\phi)$, we get $m^2 = V''(\phi_0)/[\omega(\phi_0) + 3F'^2(\phi_0)/F(\phi_0)]$. For the $f(R)$ theory, we get $m^2 = f'(0)/3f''(0)$.

In terms of the field $\tilde{h}_{\mu\nu}$,

$$
\tilde{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}\eta^{\alpha\beta}h_{\alpha\beta} - \frac{G_{4,\phi_0}}{G_4(0)}\eta_{\mu\nu}\varphi, \tag{6}
$$

and choosing the transverse traceless gauge $\partial_\mu \tilde{h}^{\mu\nu} = 0$, $\eta^{\mu\nu}\tilde{h}_{\mu\nu} = 0$ by using the freedom of coordinate transformation, the linearized Eqs. (3) and (4) become the wave equations

$$(\Box - m^2)\varphi = 0, \tag{7}$$

$$\Box \tilde{h}_{\mu\nu} = 0. \tag{8}$$

It is clear that the field $\tilde{h}_{\mu\nu}$ denotes the usual massless gravitons and it has two polarization states, the plus and cross modes. The scalar field φ is massive and it decouples from the massless tensor field $\tilde{h}_{\mu\nu}$. Suppose the massless and the massive modes both propagate in the $+z$ direction with the wave vectors,

$$k^\mu = (\Omega, 0, 0, \Omega), \quad q^\mu = (q_t, 0, 0, q_z), \tag{9}$$

respectively, where $q_t^2 - q_z^2 = m^2$. The propagation speed of the massive scalar field is $v_g = \sqrt{q_t^2 - m^2}/q_t$. The solutions to Eqs. (7) and (8) take the following form

$$\begin{aligned} \tilde{h}_{\mu\nu} &= e_{\mu\nu} e^{-ik\cdot x}, \\ \varphi &= \varphi_0 e^{-iq\cdot x}, \end{aligned} \tag{10}$$

where φ_0 and $e_{\mu\nu}$ are the amplitudes of the waves with $k^\nu e_{\nu\mu} = 0$ and $\eta^{\mu\nu} e_{\mu\nu} = 0$.

Using Eqs. (6) and (10), we get the nonvanishing Newman-Penrose variables [2, 3],

$$\Psi_4 = -\Omega^2(\tilde{h}_{xx} - i\tilde{h}_{xy}), \tag{11}$$

$$\Phi_{22} = \frac{(q_t + \sqrt{q_t^2 - m^2})^2}{4}\sigma\varphi, \tag{12}$$

$$\Phi_{00} = \frac{4(q_t - \sqrt{q_t^2 - m^2})^2}{(q_t + \sqrt{q_t^2 - m^2})^2}\Phi_{22}, \tag{13}$$

$$\Phi_{11} = -\Lambda = \frac{4m^2}{(q_t + \sqrt{q_t^2 - m^2})^2}\Phi_{22}, \tag{14}$$

where $\sigma = G_{4,\phi_0}/G_4(0)$. For different theories, we list the expressions for σ and m^2 in Table I. Due to the presence of the massive scalar field, the Newman-Penrose variables Φ_{22}, Φ_{00}, Φ_{11} and Λ do not vanish, and they are all proportional to φ. However, for the Newman-Penrose variables associated with the Weyl tensor, only Ψ_4 is nonzero and $\Psi_2 = 0$. In particular, Ψ_4 denotes the usual plus and cross modes for the massless gravitons. Note that for the massless scalar field, only Φ_{22} is nonzero and it denotes the transverse breathing mode [3].

Since the polarization states of the massless tensor field $\tilde{h}_{\mu\nu}$ are well known, we don't discuss them here. To study the polarization state due to the massive scalar mode, we need

TABLE I. The expressions for σ and m^2 in Horndeski theory, Brans-Dicke theory, the scalar-tensor theory, and $f(R)$ gravity.

	Horndeski	Brans-Dicke	Scalar-Tensor	$f(R)$
σ	$G_{4,\phi_0}/G_4(0)$	$1/\phi_0$	$F'(\phi_0)/F(\phi_0)$	$f''(0)/f'(0)$
m^2	$-\frac{K_{,\phi_0\phi_0}}{K_{,X_0}-2G_{3,\phi_0}+3G^2_{4,\phi_0}/G_4(0)}$	0	$\frac{V''(\phi_0)}{\omega(\phi_0)+3F(\phi_0)'^2/F(\phi_0)}$	$\frac{f'(0)}{3f''(0)}$

to consider the relative acceleration of freely falling test particles, i.e., we need to calculate the "electric" part of the Riemann tensor R_{tjtk} due to the massive scalar field. For the massive scalar mode, the geodesic deviation equations are,

$$\ddot{x} = -\frac{1}{2}q_t^2\sigma\varphi x, \tag{15}$$

$$\ddot{y} = -\frac{1}{2}q_t^2\sigma\varphi y, \tag{16}$$

$$\ddot{z} = -\frac{1}{2}m^2\sigma\varphi z. \tag{17}$$

Therefore, the massive scalar field φ excites both the transverse and the longitudinal polarizations, $\delta x \approx \sigma\varphi x_0/2$, $\delta y \approx \sigma\varphi y_0/2$ and $\delta z \approx (m^2/q_t^2)(\sigma\varphi/2)z_0$. Since $q_t > m$, the displacement in the transverse directions is always greater than the one in the longitudinal direction. In the massless limit, the longitudinal polarization disappears, and we get three polarization modes: breathing, plus and cross. This recovers the result in [3] for Brans-Dicke theory. In $f(R)$ gravity, we have $\sigma m^2 = 1/3$ (see Table I), so the effect of the longitudinal mode on the geodesic deviation is independent of the mass [23]. For null gravitational waves, $\Psi_2 = 0$ means that there is no longitudinal mode. However, for the massive gravitational wave, the longitudinal mode exists and the polarization state is a mixture of transverse and longitudinal mode even though $\Psi_2 = 0$.

The detection of GW150914 placed an upper bound on the mass m_g of the graviton [1],

$$m_g < 1.2 \times 10^{-22}\,\mathrm{eV}/c^2. \tag{18}$$

This potentially places severe constraint on the effect of the longitudinal polarization on the geodesic deviation in the massive case in the high frequency band, while the breathing mode can be much stronger as long as the amplitude of scalar field φ is large enough. Therefore, although there exits the longitudinal mode in the massive scalar-tensor theory, the detection of its effect is likely very difficult in the high frequency band. However, in $f(R)$ gravity, the

displacement in the longitudinal direction is independent of the mass and the displacements in the transverse directions become larger for smaller mass, so this mix mode can provide strong constraints on $f(R)$ gravity.

Table II lists the magnitudes of $|\ddot{x}^j/x^j|$ normalized with $\sigma\varphi$ for the longitudinal and the transverse modes assuming the graviton mass is given by the upper bound in Eq.(18) at three different frequencies: 100 Hz, 10^{-3} Hz, and 10^{-7} Hz. The first frequency lies in the frequency band of the ground-based detector like aLIGO and Virgo, the second frequency lies in the frequency band of space-borne observatory like LISA, and the last frequency lies in the sensitive band of pulsar timing arrays [26].

TABLE II. The dependence of the magnitudes of $|\ddot{x}^j/x^j|$ for the longitudinal and transverse modes on the frequencies of gravitational waves in units of $\mathrm{Hz}^2\sigma\varphi$.

	100 Hz	10^{-3} Hz	10^{-7} Hz
Longitudinal	1.66×10^{-14}	1.66×10^{-14}	1.66×10^{-14}
Transverse	1.97×10^{5}	1.97×10^{-5}	1.97×10^{-13}

From Table II, we see that the effect of the longitudinal mode on the geodesic deviation is smaller than that of the transverse mode by 19 to 9 orders of magnitude at higher frequencies. But at the lower frequencies, i.e., at 10^{-7} Hz, the two modes have similar amplitudes. Therefore, aLIGO/Virgo and LISA might find it difficult to detect the longitudinal mode, but pulsar time arrays should be able to detect the mix polarization state with both the longitudinal and transverse modes. The results are consistent with those for the massive graviton in a specific bimetric theory [27].

Following [14, 28], the interferometer response function of the transverse and longitudinal modes can be obtained in the frequency domain. In the interferometer, the photon is emitted from the beam splitter, bounced back by the mirror and received by the beam splitter. The propagation time when there is no gravitational wave will be different from the one when there is. For simplicity, the beam splitter is placed at the origin of the coordinate system. To obtain the response function for the longitudinal mode, project the light in the z direction

and place the mirror at $z = L$. The response function is thus given by,

$$\begin{aligned} Y_l(f) =& \frac{m^2}{16\pi^3 f^3}\Big\{4\pi f e^{i2\pi fL(v_g+1)} \\ & - \frac{e^{4i\pi fL}}{(v_g-1)^3 L}\Big[e^{i2\pi fL(v_g-1)}[2\pi fL(v_g-1) \\ & + i2] + 2\pi fL(v_g-1) - i2\Big] \\ & + \frac{1}{(v_g+1)^3 L}\Big[2\pi fL(v_g+1) + e^{i2\pi fL(v_g+1)} \\ & [2\pi fL(v_g+1) + 2i] - i2\Big]\Big\}. \end{aligned} \tag{19}$$

where $f = q_t/(2\pi)$ is the frequency of gravitational waves and the propagation speed v_g varies with the frequency. To obtain the response function for the transverse mode, project the light in the x direction and place the mirror at $x = L$. The response function is,

$$Y_b(f) = e^{i2\pi fL} + \frac{e^{i2\pi fL}[2\pi fL - \sin(2\pi fL)]}{2\pi fL} \tag{20}$$

Figure 1 shows the absolute value of the longitudinal and transverse response functions for aLIGO ($L = 4$ km) to a scalar gravitational wave with the mass $m_g = 1.2 \times 10^{-22}$ eV/c^2. Comparing the two response functions shows that the transverse response is much larger than that of the longitudinal mode in high frequencies, so the detection of the longitudinal mode becomes very difficult in the high frequency band.

III. CONCLUSION

This work analyzes the gravitational wave polarizations in the most general scalar-tensor theory of gravity, the Horndeski theory. It reveals that there are three independent polarization modes: the mixture state of the breathing and longitudinal modes for the massive scalar field, the usual plus and cross modes for the massless gravitons. These results are consistent with the three propagating degrees of freedom in Horndeski theory. Since the propagation speed of the massive gravitational wave depends on the frequency and is smaller than the speed of light, the massive mode will arrive at the detector later than the massless gravitons. Using the Newman-Penrose variables, we find that $\Psi_2 = 0$. For null gravitational waves, this means that the longitudinal mode does not exist. However, the longitudinal mode exists in the massive case even though $\Psi_2 = 0$, so the classification of the six polarization states for null gravitational waves derived from the little group $E(2)$ of the Lorentz group is not



FIG. 1. The absolute values of the longitudinal and transverse response functions $|Y_l(f)|$ and $|Y_b(f)|$ as functions of f for aLIGO ($L = 4$ km) to a scalar gravitational wave with the mass $m_g = 1.2 \times 10^{-22}$ eV/c^2. The solid curve denotes the transverse response function $|Y_b(f)|$ and the dashed curve denotes the longitudinal response function $|Y_l(f)|$.

applicable to the massive case. Although the longitudinal mode exists for the massive scalar field, it is difficult to be detected in the high frequency band because of the suppression by the extremely small graviton mass. In the massless case, the longitudinal mode disappears and the mix state reduces to the pure transverse breathing mode. We also apply the general analysis to three particular cases: Brans-Dicke theory, the general scalar-tensor theory and $f(R)$ gravity. Despite of the fact that in $f(R)$ gravity, the longitudinal mode does not depend on the mass of graviton, its magnitude is much smaller than the transverse one in the high frequency band, which makes the its detection unlikely by the network of aLIGO/VIGO and LISA, too. It is interesting that the transverse mode become stronger for smaller graviton mass in the $f(R)$ gravity, so the detection of the mixture state can place strong constraint on $f(R)$ gravity. Compared with aLIGO/VIRGO and LISA, pulsar timing arrays might be the primary tool to detect the longitudinal mode.

ACKNOWLEDGMENTS

This research was supported in part by the National Natural Science Foundation of China under Grant No. 11690021 and the Major Program of the National Natural Science

Foundation of China under Grant No. 11475065.

[1] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Phys. Rev. Lett. **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].

[2] E. Newman and R. Penrose, J. Math. Phys. **3**, 566 (1962).

[3] D. M. Eardley, D. L. Lee, and A. P. Lightman, Phys. Rev. D **8**, 3308 (1973).

[4] D. M. Eardley, D. L. Lee, A. P. Lightman, R. V. Wagoner, and C. M. Will, Phys. Rev. Lett. **30**, 884 (1973).

[5] C. Brans and R. Dicke, Phys. Rev. **124**, 925 (1961).

[6] G. W. Horndeski, Int. J. Theor. Phys. **10**, 363 (1974).

[7] M. Ostrogradsky, Mem. Acad. St. Petersbourg **6**, 385 (1850).

[8] R. Utiyama and B. S. DeWitt, J. Math. Phys. **3**, 608 (1962).

[9] K. Stelle, Phys. Rev. D **16**, 953 (1977).

[10] K. S. Stelle, Gen. Rel. Grav. **9**, 353 (1978).

[11] H. A. Buchdahl, Mon. Not. Roy. Astron. Soc. **150**, 1 (1970).

[12] J. O' Hanlon, Phys. Rev. Lett. **29**, 137 (1972).

[13] P. Teyssandier and P. Tourrenc, J. Math. Phys. **24**, 2793 (1983).

[14] C. Corda, JCAP **0704**, 009 (2007), arXiv:astro-ph/0703644 [astro-ph].

[15] C. Corda, Int. J. Mod. Phys. A **23**, 1521 (2008), arXiv:0711.4917 [gr-qc].

[16] S. Capozziello, C. Corda, and M. F. De Laurentis, Phys. Lett. B **669**, 255 (2008), arXiv:0812.2272 [astro-ph].

[17] S. Capozziello, R. Cianci, M. De Laurentis, and S. Vignolo, Eur. Phys. J. C **70**, 341 (2010), arXiv:1007.3670 [gr-qc].

[18] P. Prasia and V. C. Kuriakose, Int. J. Mod. Phys. D **23**, 1450037 (2014).

[19] M. E. S. Alves, O. D. Miranda, and J. C. N. de Araujo, Phys. Lett. B **679**, 401 (2009), arXiv:0908.0861 [gr-qc].

[20] M. E. S. Alves, O. D. Miranda, and J. C. N. de Araujo, Class. Quant. Grav. **27**, 145010 (2010), arXiv:1004.5580 [gr-qc].

[21] H. R. Kausar, L. Philippoz, and P. Jetzer, Phys. Rev. D **93**, 124071 (2016), arXiv:1606.07000 [gr-qc].

[22] Y. S. Myung, Adv. High Energy Phys. **2016**, 3901734 (2016), arXiv:1608.01764 [gr-qc].

[23] D. Liang, Y. Gong, S. Hou, and Y. Liu, "On the polarizations of gravitational waves in $f(R)$ gravity," (2017), arXiv:1701.05998 [gr-qc].

[24] W. Lin and M. Ishak, Phys. Rev. D **94**, 123011 (2016), arXiv:1605.03504 [astro-ph.CO].

[25] S. Bhattacharya and S. Chakraborty, Phys. Rev. D **95**, 044037 (2017), arXiv:1607.03693 [gr-qc].

[26] C. J. Moore, R. H. Cole, and C. P. L. Berry, Class. Quant. Grav. **32**, 015014 (2015), arXiv:1408.0740 [gr-qc].

[27] W. L. S. de Paula, O. D. Miranda, and R. M. Marinho, Class. Quant. Grav. **21**, 4595 (2004), arXiv:gr-qc/0409041 [gr-qc].

[28] M. Rakhmanov, Phys. Rev. D **71**, 084003 (2005), arXiv:gr-qc/0406009 [gr-qc].